



From Critical Moments to

Comfort

500mg & 1g I.V



Mromont is the drug of choice for severe ESBL-E (Extended Spectrum Beta-Lactamase-producing Enterobacteriaceae) infections and is used to treat bacterial meningitis, severe CAP (Community-Acquired Pneumonia), complicated UTI (urinary tract infection),

Mromont is manufactured in a dedicated, state of the art carbapenem facilit

MECHANISM OF ACTION

Meropenem is a broad-spectrum carbapenem antibiotic. It is active against Gram-positive and Gram-negative bacteria. Meropenem exerts its action by penetrating bacterial cells readily and interfering with the synthesis of vital cell wall components, which leads to cell death

Key Features and Benefits of Mromont

Broad-Spectrum Antibiotic:

Effective against a wide range of Gram-positive and Gram-negative bacteria.

Convenient Dosage:

Administered intravenously, allowing for rapid therapeutic levels.

Reasonable Price:

Offers a cost-effective solution for treating severe bacterial infections.

Good Tolerability:

Well-tolerated in both children and adults with an acceptable safety profile.

DOSAGE & ADMINISTRATION

INFECTION	ADULTS 'DOSE TO BE ADMINISTERED T.I.D.	CHILDREN FROM 3 MONTHSTO 11 YEARS OF AGE AND UP TO 50KG BODY WEIGHT T.I.D
Severe pneumonia including hospital and ventilator-associated pneumonia.	500mg or 1g	10 or 20mg/kg
Broncho-pulmonary infections in cystic fibrosis	2g	40mg/kg
Complicated urinary tract Infections	500mg or 1g	10 or 20mg/kg
Complicated intra-abdominal infections	500mg or 1g	10 or 20mg/kg
Complicated skin and soft tissue infections	500mg or 1g	10 or 20mg/kg
Acute bacterial meningitis	2g	40mg/kg
Management of febrile neutropenic patients	1g	20mg/kg
Intra and post partum infections	500mg or 1g	-

Coverage in Various Conditions:

Treats complicated skin and skin structure infections, complicated intra-abdominal infections, and bacterial meningitis

USFDA Approved for Children:

Approved for use in children aged 3 months and older with bacterial meningitis and complicated intra-abdominal infections.

High Efficacy:

Demonstrated efficacy in clinical trials for treating serious bacterial infections

Rapid Action:

Provides immediate high levels of medication in the bloodstream



